

# High-Intensity Interval Training Combined with Saffron Supplementation Modulates Stress- Inflammatory Markers in Obese Women with Type 2 Diabetes

Shadi Golpasandi<sup>1</sup>, Siamand Abdollahpour<sup>2</sup>, Hadi Golpasandi<sup>3</sup>✉

Received: 2022/01/03

Accepted: 2022/02/25

## Abstract

**Introduction:** The aim of this study was to investigate the interactive effects of saffron extract supplementation with high-intensity interval training (HIIT) on serum levels of stress-inflammatory markers in obese women with type 2 diabetes.

**Method:** Thirty-two obese women with type 2 diabetes were randomly divided into 4 intervention groups: supplement, HIIT +Supplement, placebo and placebo+ HIIT. The anthropometric and blood characteristics of subjects were evaluated in pre-test and post-test. The method of supplementation was that the supplement group received 100 mg of saffron extract daily. The subjects in the training group also did running on the treadmill in 4 bouts with intensity of 85 to 95% of the maximum heart rate along with 4 periods of active rest at the intensity of 50 to 60% of the maximum heart rate.

**Results:** Data analysis showed that the effect of time was a significant decrease in serum levels of IL-6, NT-proBNP, and GDF-15 in the supplement, supplement + HIIT+, and placebo+ HIIT groups ( $P \leq 0.05$ ). The results of time  $\times$  group interaction also showed a significant difference between the groups ( $P \leq 0.05$ ). So that the decrease in the levels of stress-inflammatory markers in the supplement + HIIT group had a greater decrease.

**Conclusion** According to the results, it can be reported that HIIT and saffron supplementation (combination of both) probably reduces the levels of stress-inflammatory markers such as GDF-15, NT-proBNP, and IL-6 by reducing fat mass, as well as also improve glucose metabolism and insulin resistance in obese women with diabetes.

**Keywords:** Growth differentiation factor-15, interleukin-6, NT-proBNP, inflammation, type 2 diabetes.

<sup>1</sup> PhD student in Medical Physiology, Department of Medical Physiology, Faculty of Medicine, Kankuk University, Seoul, South Korea.

<sup>2</sup> Master of Exercise Physiology, Department of Exercise Physiology, Faculty of Physical Education and Sport Sciences, University of Guilan, Rasht, Guilan, Iran.

<sup>3</sup> PhD student in Exercise Physiology, Department of Exercise Physiology, Faculty of Humanities and Social Sciences, University of Kurdistan, Sanandaj, Kurdistan, Iran.

✉ **Corresponding author:**  
hadigolpasandi@yahoo.com

All rights of this article are reserved for the University of Kurdistan.

## 1. Introduction

Type 2 diabetes (T2DM) is a metabolic disease that causes hyperglycemia (hyperglycemia) by impairing insulin secretion, reducing insulin sensitivity, or increasing insulin resistance(1), which can cause a reaction. Inflammatory, metabolic, and cardiovascular diseases, followed by an increased risk of disease and mortality(2). Therefore, blood glucose control is one of the cornerstones of treatment in people with type 2 diabetes(3). In addition, diabetes is one of the main risk factors for heart failure (HF), which is associated with an increased risk of more than 50% and has substantial adverse effects on HF prognosis (4). Heart failure is also the second most common symptom of cardiovascular disease (CVD) in patients with type 2 diabetes and is more common than myocardial infarction (5). However, insufficient emphasis has been placed on the prevention

and treatment of heart failure in the clinical control of diabetes (6).

Several circulating biomarkers have recently been shown to be associated with CVD, such as C-reactive protein (CRP), interleukin-6 (IL-6), and pro-BNP natriuretic peptide (NTpro-BNP), and growth differentiation factor-15 (GDF-15) (7). It has also been suggested that these biomarkers help predict the risk of heart failure. However, few studies have examined the association between these biomarkers and the risk of heart failure in diabetic patients, and it is unclear to what extent these biomarkers can classify the risk of heart failure in such patients.

In response, traditional and emerging medication strategies and lifestyles such as physical activity and diet continue to be considered to reduce the burden of these metabolic disease conditions (8, 9). The American

## Citation:

Golpasandi, Sh., Abdollahpour, S., Golpasandi, H. High-intensity interval training combined with saffron supplementation modulates stress-inflammatory markers in obese women with type 2 diabetes. *Research in Exercise Nutrition*, 2022; 1(1): 61-55. doi: 10.34785/J019.2022.002.

Diabetes Association (ADA) recommends at least 150 minutes a week of moderate to vigorous physical activity or 75 minutes a week of vigorous physical activity for everyone (8, 9). However, inactivity and lack of perceived time are the main reported reasons for not exercising regularly (10, 11). Many people today use new exercise training regimens such as high-intensity interval training (HIIT) with active intervals (30 seconds and 4 minutes) with active rest intervals. On the other hand, the cost-effectiveness of this type of exercise in terms of time, as well as similar metabolic and physiological adaptations (12-14) and sometimes superior (15, 16) of this type of exercise compared to continuous moderate-intensity training (MICT) causes it has become increasingly popular.

On the other hand, nowadays, they are active food compounds or dietary supplements that have been proven to have benefits beyond basic nutritional functions, including health promotion through treatment and prevention of diseases (17), which is a factor in various components of plants such as flowers, There are roots in plants and oils called "nutrients" (18). Saffron is a short-lived plant that is found in various parts of the Mediterranean, Europe, and Asia (19). Saffron is used as a spice due to its coloring, flavoring, and aromatic properties (20). *Crocus sativus* contains crocetin, crocin, pyrocrocetin, and safranal which are effective in regulating inflammation and oxidative stress (21). In line with the anti-inflammatory properties of saffron, it can also be said that saffron prevents the activation and release of kappa nuclear factor activated B-cell light chain (NF- $\kappa$ B) and prevents the transfer of NF- $\kappa$ B to the nucleus following the transcription of inflammation-related genes. However, it can be said that saffron exerts its anti-inflammatory properties by reducing the production of inflammatory proteins such as TNF- $\alpha$ , interleukin-1 $\beta$  (IL-1 $\beta$ ), and IL-6 (22). Contrary to the generally favorable relationship between saffron supplementation and subclinical inflammation indices found in laboratory and animal studies, conflicting results have been reported from clinical trials, although some studies have reported a beneficial effect of saffron consumption on inflammatory cytokines. But others could not show a significant effect (23-26).

Therefore, while saffron supplementation seems to have beneficial effects in preventing the complications of diabetes, its role in factor with exercise training on some indicators of stress and inflammation is unclear. The present study aimed to investigate the interactive effect of 8 weeks of high-intensity intermittent training with saffron supplementation on cardiac stress markers and glycemic control in obese women with type 2 diabetes.

## 2. Methods

The present research design is quasi-experimental. The study population consisted of adult women aged 40 to 60 years, who after being summoned to the Diabetes Clinic and Medical Centers of Urmia, 35 people

announced their presence. Criteria for inclusion in the present study include no history of regular exercise training, body mass index (BMI) of more than 30 kg/m<sup>2</sup>, high body fasting blood sugar level above 130 mg/dl, no history of smoking, and no history of cardiovascular disease. After evaluating the inclusion criteria, 32 subjects were divided into 4 groups: 1. Saffron, 2. Saffron + HIIT, 3. Placebo, and 4. Placebo + HIIT.

In the initial evaluation session, the subjects were introduced to how to implement and apply the research interventions and then completed the questionnaires related to medical health and voluntary participation in the research. In the next session, pre-test anthropometric indices, including weight, height, body mass index (BMI), waist-to-hip ratio (WHR), and body fat percentage, also cardiac stress markers including IL-6, NT-proBNP, and GDF-15 were taken from the brachial vein blood sample (5 cc).

### Saffron supplementation protocol

Subjects consumed a powder capsule containing 100 mg of saffron and starch daily under the interventions, supplement + HIIT training, placebo, and placebo + HIIT, respectively (27).

### Measurement of maximum oxygen consumption (VO<sub>2</sub>max) and training protocol.

Bruce running test was used to measure maximum oxygen consumption (VO<sub>2</sub>max). This test is consisting of 7 steps three-minute and starts at a speed of 1.7 km/h with a 10% slope on the treadmill. After 10 minutes of warm-up, subjects started the test. After voluntary fatigue (according to the pressure perception criterion), the test was stopped, and their heart rate was recorded at the point of exhaustion (28). The primary training protocol also included: 8 weeks, 3 sessions per week, 4 running bouts on a treadmill with an intensity of 85-95% of maximum heart rate (MHR), and 4 rest interval periods with an intensity of 50-60% (MHR) (29). 72 hours after the last training protocol session, anthropometric characteristics and blood serum indices (stress markers) were measured using the ELISA method. Blood samples were taken immediately to separate serum from blood plasma using a centrifuge for 15 minutes at a speed of 3 meters per minute.

### Statistical Analysis

The statistical method of the present study was performed using descriptive statistics indicators, including mean and standard deviation in the form of tables and graphs using Excel software. The research findings were analyzed using a two-way analysis of variance (4 \* 2) using SPSS software version 25 at the level ( $P \leq 0.05$ ).

## 3. Results

Mean, and standard deviation of descriptive, anthropometric characteristics including age, height, weight, BMI, fasting glucose and insulin levels and glycosylated hemoglobin (HbA1c) are shown in Table 1.

**Table (1): Mean and standard deviation (M ± SD) of descriptive and anthropometric indices of the subjects**

Groups	Saffron		Saffron + HIIT		Placebo		Placebo + HIIT	
	Pre- test	Post-test	pre- test	Post-test	pre- test	Post-test	pre- test	Post-test
<b>Intervention times</b>								
<b>Age (years)</b>	47/48 ± 7/68		49/18 ± 7/34		47/36 ± 8/29		45/05 ± 9/10	
<b>Height (cm)</b>	162 ± 8/84		161/12 ± 5/78		159/25 ± 6/53		160/35 ± 6/23	
<b>Weight (kg)</b>	80/88 ± 10/37	79/49 ± 11/74	82/10 ± 9/31	78/52 ± 8/43	81/37 ± 10/02	81/50 ± 8/55	80/13 ± 11/74	79/52 ± 9/39
<b>BMI (Kg/m2)</b>	30/35 ± 2/50	29/35 ± 2/45	32/23 ± 2/02	29/14 ± 1/69	32/71 ± 2/15	32/32 ± 2/60	31/78 ± 1/41	30/64 ± 1/18
<b>Fasting blood Glucose (ml/dL)</b>	165/25 ± 43/6	160/15 ± 39/19	167/68 ± 48/8	158/26 ± 61/24	165/09 ± 39/28	164/11 ± 40/06	160/21 ± 46/1	157/90 ± 42/25
<b>Fasting Blood insulin (pmol/L)</b>	160/34 ± 23/38	156/51 ± 18/36	155/51 ± 18/65	149/13 ± 19/41	161/28 ± 28/12	160/08 ± 26/19	153/59 ± 21/34	151/26 ± 20/78
<b>HbA1c</b>	7/54 ± 1/67	7/13 ± 1/04	7/31 ± 1/48	6/67 ± 1/08	7/23 ± 1/43	7/18 ± 1/18	6/89 ± 1/78	6/17 ± 1/27

The results of IL-6, NT-proBNP, and GDF-15 levels are shown in Table 2 and Figure 1, respectively.

**Table (2): Mean and standard deviation of IL-6 and NT-proBNP levels among research groups**

Variables	Saffron		Saffron + HIIT		Placebo		Placebo + HIIT	
	Pre-test	Post-test	Pre-test	Post-test	Pre-test	Post-test	Pre-test	Post-test
<b>IL-6 (pg/ml)</b>	90.16 ± 15.04	89.37 ± 16.18	89.36 ± 13.08	86.50 ± 14.04 *	90.68 ± 14.11	90.13 ± 16.68	89.57 ± 15.28	88.31 ± 17.14
<b>NT-proBNP (pg/ml)</b>	651.34 ± 119.18	618.87 ± 121.23 *	648.35 ± 100.44	598.64 ± 115.29 *¥	678.41 ± 100.12	669.16 ± 95.14	699.16 ± 106.13	686.73 ± 116.35 *

\* Significance relative to pre-test time ( $P \leq 0.05$ ). ¥ Significance relative to groups ( $P \leq 0.05$ ).

+ HIIT groups in the post-test compared to pre-test ( $P \leq 0.000$ ) so that their levels are significantly reduced. However, this decrease was greater among the HIIT + Saffron group than in the other groups ( $P \leq 0.001$ ).

The results of the analysis of variance ( $4 \times 2$ ) test related to IL-6 and NT-proBNP variables in Table 2 show a significant difference between the levels of both components in the Saffron, Saffron + HIIT, and placebo




Figure (1): Changes in serum GDF-15 levels in the subjects of the present study.

\* Significance relative to pre-test time ( $P \leq 0.05$ ). ¥ Significance relative to groups ( $P \leq 0.05$ ).

According to the results (Figure 1) related to changes in GDF-15 levels, it can be said that GDF-15 levels in the Saffron, Saffron + HIIT, and placebo + HIIT groups in the post-test decreased significantly compared to the pre-test ( $P \leq 0.000$ ). However, this decrease is more significant in the HIIT + Saffron group ( $P \leq 0.003$ ).

#### 4. Discussion

This study aimed to investigate the interactive effect of 8 weeks of HIIT training with saffron supplementation on stress and inflammatory indices in obese women with type 2 diabetes. The results showed that IL-6 and NT-proBNP levels in the Saffron, Saffron + HIIT, and placebo + HIIT groups were significantly reduced in the post-test compared to the pre-test (time effect). These results were consistent with the findings of Silva et al.

(2022), Francois et al. (2015), Liu et al. (2019), and Delfan et al. (2020) (30-33). In previous studies, stress indicators in response to exercise have not been specifically studied. However, the study of critical cardiac markers in people with type 2 diabetes as one of the predictors of heart failure was shown that higher levels of IL-6, hs-CRP, hs-cTnT, and NT-proBNP were significantly associated with an increased risk or progression of heart failure in patients with type 2 diabetes (7). Type 2 diabetes is characterized by more insulin resistance in skeletal muscle and a lack of insulin secretion in the final stage. In general, T2DM increases blood glucose levels and components of the metabolic syndrome, including hypertension, triglycerides, decreased high-density lipoprotein levels, and abdominal obesity (34). Increased inflammatory factors including tumor necrosis factor (TNF- $\alpha$ ), reactive protein C (CRP), and IL-6 due to increased abdominal obesity, different signals to regulate various factors including insulin sensitivity, stress Sends oxidative, energy metabolism, blood clotting and inflammatory responses (35). GDF-15 has recently been identified as an important plasma marker associated with cardiac metabolic syndrome. GDF-15 has been termed the macrophage inhibitor cytokine-1 (MIC-1) due to its inhibition of TNF production in lipopolysaccharide-stimulated macrophages(36). Recently, GDF-15 has been shown to reduce food intake, body weight, and obesity and improve glucose tolerance in typical, high-fat diets (37). In a previous study, it was found that acute exercise significantly increased serum GDF-15 levels in obese men, possibly improving glucose metabolism in them, whereas levels decreased significantly after a few hours(38). In the present study, it was also shown that eight weeks of high-intensity interval training along with saffron supplementation caused significant levels. However, it can be said that HIIT training along with saffron supplementation due to their anti-inflammatory properties reduce serum GDF-15 levels in obese women with type 2 diabetes, which these changes were associated with improved anthropometric indices such as Weight and BMI. Higher levels of GDF-15 and a known cardiovascular risk factor such as NT-proBNP have also been a predictor of cardiovascular mortality in patients with diabetic nephropathy (39). However, there are limited studies on the effect of exercise on GDF-15 and NT-proBNP marker levels. Ebadi et al. (2021) reported that eight weeks of HIIT training with astaxanthin supplementation significantly affected NT-proBNP levels and insulin resistance in diabetic rats (40). In another study, Galliera et al. (2014) showed that GDF-15 and NT-proBNP levels significantly increased in rugby players after strenuous exercise sessions (41). This study emphasized the role of two essential factors, GDF-15 and NTproBNP as markers for evaluating cardiovascular function in players. However, Munk et al. (2014) found that long-term exercise training did not affect GDF-15 and NT-proBNP levels (42), which was inconsistent with the present study results. One of the reasons for this discrepancy was probably the type of subjects, so they studied people with coronary artery

disease, while in the present study, the subjects were obese women with type 2 diabetes.

The biologically active components of saffron have been reported to exert responses to inflammatory and anti-inflammatory agents through their effect on insulin resistance. According to previous reports, crocin, as an active ingredient in saffron, can inhibit the production of IL-6 and TNF- $\alpha$ , thereby activating anti-inflammatory signals in hyperglycemic conditions (43). In addition, it is suggested that the anti-inflammatory activity of saffron is due to its potent antioxidant properties and its radical inhibition. Saffron may suppress the increase in oxidative stress induced in type 2 diabetes and may begin by depleting reactive oxygen species, strengthening the antioxidant defense system, and regulating the immune response (44). In our study, supplementation of saffron as a 100 mg capsule per day for eight weeks reduced the inflammatory marker IL-6, which was observed in both the supplement group and the supplement group + HIIT training. However, these changes were more significant in the combined group (supplement + HIIT training), probably due to the anti-inflammatory properties of saffron and HIIT training. One of the mechanisms through which exercise training reduces the serum levels of inflammatory cytokines, including IL-6, is the reduction of fat mass and the expression of inflammatory mediators caused by that (45), in the present study, these changes were associated with a decrease in fat mass and BMI subjects.

Exercise volume and intensity as the basis of exercise training protocol is another important factor in controlling the risk of diabetes and its complications. Although low- or moderate-intensity continuous aerobic training is recommended for use in diabetics, high-intensity intermittent training (HIIT) has been reported to be more effective in rejuvenating the heart, controlling blood glucose, and many other clinical complications of diabetes (32). Although low- or moderate-intensity continuous aerobic training is recommended for use in diabetics, high-intensity intermittent training (HIIT) has been reported to be more effective in optimal heart function, blood glucose control, and many other clinical complications diabetes (32). Støa et al. (2017) found that the HIIT training program with an intensity of 85-95% of maximum heart rate significantly increased  $VO_{2peak}$  and decreased hemoglobin A1c (HbA1c), body weight and eight in subjects with T2D (46). Karstoft et al. (2013) observed the effectiveness of HIIT in further improvements in  $VO_{2peak}$ , body weight, fat mass, and glycemic control compared with continuous training in the form of walking (47). Mitranun et al. (2014), also reported that HIIT improved cardiovascular risk factors, including HbA1c, glycemic control, and  $Vo_{2max}$ , as well as other cardiovascular risk factors in T2D patients compared to traditional continuous training (48), which was consistent with the results of the present study.

In conclusion, the results of the present study showed that HIIT training and saffron supplementation (combination of both) possibly improved glucose metabolism through anti-inflammatory properties due to the reduction of levels of stress-inflammatory markers



such as GDF-15, NT-proBNP, and IL-6 improving glucose metabolism and insulin resistance in obese women with type 2 diabetes.

**Acknowledgments:** We would like to thank the participants for contribution to the study.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

- [1] Rabinovitch A, Suarez-Pinzon WL. Cytokines and their roles in pancreatic islet  $\beta$ -cell destruction and insulin-dependent diabetes mellitus. *Biochemical pharmacology*. 1998;55(8):1139-49.
- [2] Jiménez-Maldonado A, García-Suárez PC, Rentería I, Moncada-Jiménez J, Plaisance EP. Impact of high-intensity interval training and sprint interval training on peripheral markers of glycemic control in metabolic syndrome and type 2 diabetes. *Biochimica et Biophysica Acta (BBA)-Molecular Basis of Disease*. 2020;1866(8):165820.
- [3] Nathan DM, Buse JB, Davidson MB, Ferrannini E, Holman RR, Sherwin R, et al. Medical management of hyperglycemia in type 2 diabetes: a consensus algorithm for the initiation and adjustment of therapy: a consensus statement of the American Diabetes Association and the European Association for the Study of Diabetes. *Diabetes care*. 2009;32(1):193-203.
- [4] Pocock SJ, Wang D, Pfeffer MA, Yusuf S, McMurray JJ, Swedberg KB, et al. Predictors of mortality and morbidity in patients with chronic heart failure. *European heart journal*. 2006;27(1):65-75.
- [5] Shah AD, Langenberg C, Rapsomaniki E, Denaxas S, Pujades-Rodriguez M, Gale CP, et al. Type 2 diabetes and incidence of cardiovascular diseases: a cohort study in 1· 9 million people. *The lancet Diabetes & endocrinology*. 2015;3(2):105-13.
- [6] McMurray JJ, Gerstein HC, Holman RR, Pfeffer MA. Heart failure: a cardiovascular outcome in diabetes that can no longer be ignored. *The lancet Diabetes & endocrinology*. 2014;2(10):843-51.
- [7] Ohkuma T, Jun M, Woodward M, Zoungas S, Cooper ME, Grobbee DE, et al. Cardiac stress and inflammatory markers as predictors of heart failure in patients with type 2 diabetes: the ADVANCE trial. *Diabetes Care*. 2017;40(9):1203-9.
- [8] Association AD. 13. Children and adolescents: standards of medical care in diabetes—2021. *Diabetes care*. 2021;44(Supplement\_1):S180-S99.
- [9] Colberg SR, Sigal RJ, Yardley JE, Riddell MC, Dunstan DW, Dempsey PC, et al. Physical activity/exercise and diabetes: a position statement of the American Diabetes Association. *Diabetes care*. 2016;39(11):2065-79.
- [10] Chang C, Khurana S, Strodel R, Camp A, Magenheimer E, Hawley N. Perceived barriers to physical activity among low-income Latina women at risk for type 2 diabetes. *The Diabetes Educator*. 2018;44(5):444-53.
- [11] Thomas N, Alder E, Leese G. Barriers to physical activity in patients with diabetes. *Postgraduate medical journal*. 2004;80(943):287-91.
- [12] Burgomaster KA, Howarth KR, Phillips SM, Rakobowchuk M, MacDonald MJ, McGee SL, et al. Similar metabolic adaptations during exercise after low volume sprint interval and traditional endurance training in humans. *The Journal of physiology*. 2008;586(1):151-60.
- [13] Gibala MJ, Little JP, Van Essen M, Wilkin GP, Burgomaster KA, Safdar A, et al. Short-term sprint interval versus traditional endurance training: similar initial adaptations in human skeletal muscle and exercise performance. *The Journal of physiology*. 2006;575(3):901-11.
- [14] Heiskanen MA, Leskinen T, Heinonen IH, Löyttyniemi E, Eskelinen J-J, Virtanen K, et al. Right ventricular metabolic adaptations to high-intensity interval and moderate-intensity continuous training in healthy middle-aged men. *American Journal of Physiology-Heart and Circulatory Physiology*. 2016;311(3):H667-H75.
- [15] MacInnis MJ, Gibala MJ. Physiological adaptations to interval training and the role of exercise intensity. *The Journal of physiology*. 2017;595(9):2915-30.
- [16] Chavanelle V, Boisseau N, Otero YF, Combaret L, Dardevet D, Montaurier C, et al. Effects of high-intensity interval training and moderate-intensity continuous training on glycaemic control and skeletal muscle mitochondrial function in db/db mice. *Scientific reports*. 2017;7(1):1-10.
- [17] Weaver CM. Bioactive foods and ingredients for health. *Advances in Nutrition*. 2014;5(3):306S-11S.
- [18] Ghaffari S, Roshanravan N. Saffron; An updated review on biological properties with special focus on cardiovascular effects. *Biomedicine & Pharmacotherapy*. 2019;109:21-7.
- [19] Ghaffari S, Roshanravan N. The role of nutraceuticals in prevention and treatment of hypertension: An updated review of the literature. *Food Research International*. 2020;128:108749.
- [20] Jose Bagur M, Alonso Salinas GL, Jiménez-Monreal AM, Chaouqi S, Llorens S, Martínez-Tomé M, et al. Saffron: An old medicinal plant and a potential novel functional food. *Molecules*. 2017;23(1):30.
- [21] Khatir SA, Bayatian A, Barzegari A, Roshanravan N, Safaiyan A, Pavon-Djavid G, et al. Saffron (*Crocus sativus* L.) supplements modulate circulating MicroRNA (miR-21) in atherosclerosis patients; a randomized, double-blind, placebo-controlled trial. *Iranian Red Crescent Medical Journal*. 2018;20(10).
- [22] Cai J, Yi FF, Bian ZY, Shen DF, Yang L, Yan L, et al. Crocetin protects against cardiac hypertrophy by blocking MEK-ERK1/2 signalling pathway. *Journal of cellular and molecular medicine*. 2009;13(5):909-25.

- [23] Hazman Ö, Ovalı S. Investigation of the anti-inflammatory effects of safranal on high-fat diet and multiple low-dose streptozotocin induced type 2 diabetes rat model. *Inflammation*. 2015;38(3):1012-9.
- [24] Azimi P, Ghiasvand R, Feizi A, Hariri M, Abbasi B. Effects of cinnamon, cardamom, saffron, and ginger consumption on markers of glycemic control, lipid profile, oxidative stress, and inflammation in type 2 diabetes patients. *The review of diabetic studies: RDS*. 2014;11(3):258.
- [25] Christodoulou E, Kadoglou N, Stasinopoulou M, Konstandi O, Kenoutis C, Kakazanis Z, et al. *Crocus sativus* L. aqueous extract reduces atherogenesis, increases atherosclerotic plaque stability and improves glucose control in diabetic atherosclerotic animals. *Atherosclerosis*. 2018;268:207-14.
- [26] Ebrahimi F, Sahebkar A, Aryaeian N, Pahlavani N, Fallah S, Moradi N, et al. Effects of saffron supplementation on inflammation and metabolic responses in type 2 diabetic patients: A randomized, double-blind, placebo-controlled trial. *Diabetes, metabolic syndrome and obesity: targets and therapy*. 2019;12:2107.
- [27] Mobasser M, Ostadrahimi A, Tajaddini A, Asghari S, Barati M, Akbarzadeh M, et al. Effects of saffron supplementation on glycemia and inflammation in patients with type 2 diabetes mellitus: a randomized double-blind, placebo-controlled clinical trial study. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*. 2020;14(4):527-34.
- [28] Vittal V, Shankar V, Kutty K, Kumar SA. An Interventional Study of Physical Activity on HbA1C and VO<sub>2</sub>max in Type 2 Diabetes Mellitus. *Indian journal of physiology and pharmacology*. 2016;60(1):76-81.
- [29] Ahmad AM. Moderate-intensity continuous training: is it as good as high-intensity interval training for glycemic control in type 2 diabetes? *Journal of exercise rehabilitation*. 2019;15(2):327.
- [30] Silva LRB, Gentil P, Seguro CS, de Oliveira JCM, Silva MS, Marques VA, et al. High-Intensity Interval Training Improves Cardiac Autonomic Function in Patients with Type 2 Diabetes: A Randomized Controlled Trial. *Biology*. 2022;11(1):66.
- [31] Francois ME, Little JP. Effectiveness and safety of high-intensity interval training in patients with type 2 diabetes. *Diabetes Spectrum*. 2015;28(1):39-44.
- [32] Delfan M, Delphan M, Kordi MR, Ravasi AA, Safa M, Gorgani-Firuzjaee S, et al. High intensity interval training improves diabetic cardiomyopathy via miR-1 dependent suppression of cardiomyocyte apoptosis in diabetic rats. *Journal of Diabetes & Metabolic Disorders*. 2020;19(1):145-52.
- [33] Liu J-x, Zhu L, Li P-j, Li N, Xu Y-b. Effectiveness of high-intensity interval training on glycemic control and cardiorespiratory fitness in patients with type 2 diabetes: a systematic review and meta-analysis. *Aging clinical and experimental research*. 2019;31(5):575-93.
- [34] Kaur J. A comprehensive review on metabolic syndrome. *Cardiology research and practice*. 2014;2014.
- [35] Sivitz WI. Lipotoxicity and glucotoxicity in type 2 diabetes: effects on development and progression. *Postgraduate medicine*. 2001;109(4):55-64.
- [36] Adela R, Banerjee SK. GDF-15 as a target and biomarker for diabetes and cardiovascular diseases: a translational prospective. *Journal of diabetes research*. 2015;2015.
- [37] Macia L, Tsai VW-W, Nguyen AD, Johnen H, Kuffner T, Shi Y-C, et al. Macrophage inhibitory cytokine 1 (MIC-1/GDF15) decreases food intake, body weight and improves glucose tolerance in mice on normal & obesogenic diets. *PloS one*. 2012;7(4):e34868.
- [38] Golpasandi, H., Mirzaei, B., & Golpasandi, S. (2019). Glucose, Insulin and Growth/Differentiation Factor-15 Serums Responses to the Acute Effects of Two Intermittent and Continuous Exercises in Sedentary Obese Males. *Journal of Inflammatory Diseases*, 23(4), 286-295.
- [39] Van Huyen JPD, Cheval L, Bloch-Faure M, Belair MF, Heudes D, Bruneval P, et al. GDF15 triggers homeostatic proliferation of acid-secreting collecting duct cells. *Journal of the American Society of Nephrology*. 2008;19(10):1965-74.
- [40] Ebadi, N., Zolfaghari, M. R., & Ghaderi Pakdel, F. (2021). The effect of aerobic and interval training with the astaxanthin supplementation on cardiac biomarkers and insulin resistance in type 2 diabetic rats. *Daneshvar Medicine*, 28(5), 23-36.
- [41] Galliera E, Lombardi G, Marazzi MG, Grasso D, Vianello E, Pozzoni R, et al. Acute exercise in elite rugby players increases the circulating level of the cardiovascular biomarker GDF-15. *Scandinavian Journal of Clinical and Laboratory Investigation*. 2014;74(6):492-9.
- [42] Munk PS, Valborgland T, Butt N, Larsen AI. Response of growth differentiation factor-15 to percutaneous coronary intervention and regular exercise training. *Scandinavian Cardiovascular Journal*. 2011;45(1):27-32.
- [43] Samarghandian S, Azimi-Nezhad M, Farkhondeh T. Crocin attenuate Tumor Necrosis Factor-alpha (TNF- $\alpha$ ) and interleukin-6 (IL-6) in streptozotocin-induced diabetic rat aorta. *Cytokine*. 2016;88:20-8.
- [44] Boskabady MH, Farkhondeh T. Antiinflammatory, antioxidant, and immunomodulatory effects of *Crocus sativus* L. and its main constituents. *Phytotherapy Research*. 2016;30(7):1072-94.
- [45] Karstoft K, Pedersen BK. Exercise and type 2 diabetes: focus on metabolism and inflammation. *Immunology and cell biology*. 2016;94(2):146-50.
- [46] Støa EM, Meling S, Nyhus L-K, Strømstad G, Mangerud KM, Helgerud J, et al. High-intensity aerobic interval training improves aerobic fitness and HbA1c among persons diagnosed with type 2 diabetes. *European journal of applied physiology*. 2017;117(3):455-67.

- [47] Karstoft K, Winding K, Knudsen SH, Nielsen JS, Thomsen C, Pedersen BK, et al. The effects of free-living interval-walking training on glycemic control, body composition, and physical fitness in type 2 diabetic patients: a randomized, controlled trial. *Diabetes care*. 2013;36(2):228-36.
- [48] Mitranun W, Deerochanawong C, Tanaka H, Suksom D. Continuous vs interval training on glycemic control and macro-and microvascular reactivity in type 2 diabetic patients. *Scandinavian journal of medicine & science in sports*. 2014;24(2):e69-e76.